CONSTRAINING TOPOLOGY WITH THE CMB

Angélica de Oliveira-Costa[1], George F. Smoot[2] & Alexei A. Starobinsky[3]

[1] *Max-Planck-Institut für Astrophysik, Garching, Germany.*

[2] *Lawrence Berkeley National Laboratory, Berkeley, USA.*

[3] *Russian Academy of Sciences, Moscow, Russia.*

Abstract

We present a new data analysis method to study rectangular T^3 "small universes" with one or two of its dimensions significantly smaller than the present horizon (which we refer to as T^1- and T^2-models, respectively). We find that the 4 year *COBE*/DMR data set a lower limit on the smallest cell size for T^1- and T^2-models of $3000h^{-1}$Mpc, at 95% confidence, for a scale invariant power spectrum (n=1).

1 Introduction

In the past few years, mainly after the discovery of CMB anisotropies by *COBE*/DMR [11], the study of the topology of the universe has become an important problem for cosmologists and some hypotheses, such as the "small universe" model [4], have received considerable attention. From the theoretical point of view, it is possible to have quantum creation of the universe with a multiply-connected topology [16]. From the observational side, this model has been used to explain the "observed" periodicity in the distributions of quasars [7] and galaxies [1].

Almost all work on "small universes" has been limited to the case where the spatial sections form a rectangular basic cell with sides L_x, L_y, L_z and with opposite faces topologically connected, a topology known as toroidal. The three-dimensional cubic torus T^3 is the simplest model among all possible multiply-connected topologies, in which all three sides have the same size $L \equiv L_x = L_y = L_z$. In spite of the fact that cubic T^3-model has beenΘ ruled out by *COBE* results [2, 8, 12, 13, 14], the possibility that we live in a universe with a more anisotropic topology, such as a rectangular torus T^3, is an open option that has not been ruled out yet. For instance, if the toroidal model is not a cube, but a rectangle with sides $L_x \neq L_y \neq L_z$ and with one or two of its dimensions significantly smaller than the horizon R_H ($\equiv 2cH_0^{-1}$), this small rectangular universe cannot be completely excluded by any of the previous analyses:



Figure 1: Simulated sky maps for the T^1- and T^2-models and their S-maps. (A) T^1-model with dimensions $(L_x, L_y, L_z) = R_H(3{,}3{,}0.3)$; (B) T^2-model with dimensions $(L_x, L_y, L_z) = R_H(0.3{,}0.3{,}3)$; (C) S-map of the T^1-model shown in (A); (D) S-map of the T^2-model shown in (B). Both models are plotted in galactic coordinates and have a scale invariant power spectrum (n=1).

constraints from the DMR data merely require that at least one of the sides of the cell be larger than R_H.

As pointed out by [5] and [13], if the rectangular T^3-universe has one of the cell sizes smaller than the horizon and the other two cell sizes are of the order of or larger than the horizon (for instance, $L_z \ll R_H$ and $L_x, L_y \gtrsim R_H$), the values of $\delta T/T$ are almost independent of the z-coordinate, *i.e.*, the large scale CMB pattern shows the existence of a symmetry plane formed by the x and y-axes; and if it has two cell sizes smaller than the horizon and the third cell size is of the order of or larger than the horizon (for instance, $L_x, L_y \ll R_H$ and $L_z \gtrsim R_H$), the temperatures $\delta T/T$ are aproximately independent of both x and y, *i.e.*, the CMB pattern shows the existence of a symmetry axis: values of $\delta T/T$ are almost constant along rings around the z-axis. We call the former case a T^1-model because the spatial topology of the universe becomes just T^1 in the limit $L_x, L_y \to \infty$ with L_z being fixed. The later case is denoted a T^2-model for the same reason (the corresponding limit is $L_z \to \infty$ with L_x, L_y being fixed). See Figures 1A (upper left) and 1B (upper right).

Our goal is to show that the $COBE$/DMR maps have the ability to test and rule out T^1- and T^2-models. We use a new approach to study these models in which we constrain their sizes by looking for the symmetries that they would produce in the CMB, obtaining strong constraints from the 4 year $COBE$/DMR data.

2 The method

The analysis of T^1- and T^2-models is not an easy task, since there are infinitely many combinations of different cell sizes and cell orientations. In order to study these models, we choose a statistic in which we calculate the function $S(\hat{\mathbf{n}}_i)$ defined by [3]

$$S(\hat{\mathbf{n}}_i) \equiv \frac{1}{N_{pix}} \sum_{j=1}^{N_{pix}} \frac{[\frac{\delta T}{T}(\hat{\mathbf{n}}_j) - \frac{\delta T}{T}(\hat{\mathbf{n}}_{ij})]^2}{\sigma(\hat{\mathbf{n}}_j)^2 + \sigma(\hat{\mathbf{n}}_{ij})^2}, \tag{1}$$

where N_{pix} is the number of pixels that remain in the map after the Galaxy cut has taken place, $\hat{\mathbf{n}}_{ij}$ denotes the reflection of $\hat{\mathbf{n}}_j$ in the plane whose normal is $\hat{\mathbf{n}}_i$, *i.e.*,

$$\hat{\mathbf{n}}_{ij} = \hat{\mathbf{n}}_j - 2(\hat{\mathbf{n}}_i \cdot \hat{\mathbf{n}}_j)\hat{\mathbf{n}}_i, \tag{2}$$

and $\sigma(\hat{\mathbf{n}}_j)$ and $\sigma(\hat{\mathbf{n}}_{ij})$ are the r.m.s. errors associated with the pixels in the directions $\hat{\mathbf{n}}_j$ and $\hat{\mathbf{n}}_{ij}$. $S(\hat{\mathbf{n}}_i)$ is a measure of how much reflection symmetry there is in the mirror plane perpendicular to $\hat{\mathbf{n}}_i$. The more perfect the symmetry is, the smaller $S(\hat{\mathbf{n}}_i)$ will be. When we calculate $S(\hat{\mathbf{n}}_i)$ for all 6144 pixels at the positions $\hat{\mathbf{n}}_i$, we obtain a sky map that we refer to as an S-map. This sky map is a useful visualization tool and gives intuitive understanding of how the statistic $S(\hat{\mathbf{n}}_i)$ works.

In order to better understand $S(\hat{\mathbf{n}}_i)$, we first consider the simple model of a T^1-universe with $L_z \ll R_H$. For this specific model, the values of $\delta T/T$ are almost independent of the z-coordinate, so we have almost perfect mirror symmetry about the xy-plane or, in spherical coordinates, $\delta T/T(\theta,\phi) \approx \delta T/T(\pi-\theta,\phi)$. When $\hat{\mathbf{n}}_i$ points in the direction of the smallest cell size (*i.e.*, in the z-direction), we have $S(\hat{\mathbf{n}}_i) \approx 1$; otherwise, $S(\hat{\mathbf{n}}_i) > 1$. An S-map for a T^1-model $(L_x, L_y, L_z) = R_H$(3,3,0.3) can be seen in Figure 1C (lower left). Notice in this plot that the direction in which the cell is smallest can be easily identified by two "dark spots" at $\hat{\mathbf{n}}_i \approx \hat{\mathbf{z}}$ and $\hat{\mathbf{n}}_i \approx -\hat{\mathbf{z}}$. For T^2-models, the only difference will be that in the place of the two "dark spots", we have a "dark ring" structure in the plane formed by the two small directions. See Figure 1D (lower right), an S-map of the T^2-model $(L_x, L_y, L_z) = R_H$(0.3,0.3,3).

From the two S-maps, we can infer two important properties: first, the direction in which the S-map takes its minimum value, denoted $S_\circ$, is the direction in which the universe is small. For a large universe such as $L_x, L_y, L_z \gg R_H$, the $S_\circ$-directions obtained from different realizations are randomly distributed in the sky. Secondly, the distribution of $S_\circ$-values changes with the cell size, *i.e.*, as the universe becomes smaller, the values of $S_\circ$ decrease. From the definition of the S-map, it is easy to see that the value of $S_\circ$ is independent of the cell orientation. In other words, if we rotate the cell, we will just be rotating the S-map, leaving its minimum value $S_\circ$ unchanged.

From here on, we will present our results in terms of the cell sizes R_x, R_y and R_z, usually sorted as $R_x \leq R_y \leq R_z$ and defined as $R_x \equiv L_x/R_H$, $R_y \equiv L_y/R_H$ and $R_z \equiv L_z/R_H$. We remind the reader that the results are identical for all six permutations of R_x, R_y and R_z.

3 Data Analysis

If the density fluctuations are adiabatic and the Universe is spatially flat, the Sachs-Wolfe fluctuations in the CMB are given by [10]

$$\frac{\delta T}{T}(\theta,\phi) = -\frac{1}{2}\frac{H_0^2}{c^2}\sum_{\mathbf{k}} \frac{\delta_{\mathbf{k}}}{k^2} e^{i\mathbf{k}\cdot\mathbf{r}}, \tag{3}$$

where $\mathbf{r}$ is the vector with length $R_H \equiv 2cH_0^{-1}$ that points in the direction of observation (θ,ϕ), H_0 is the Hubble constant (written here as $100h$ km s^{-1} Mpc^{-1}) and $\delta_{\mathbf{k}}$ is the density fluctuation in Fourier space, with the sum taken over all wave numbers $\mathbf{k}$.



Figure 2: Cumulative probability distribution of $S_\circ$ for T^1- and T^2-models obtained from Monte Carlo simulations. A, left plot: Simulations for T^2-universes with dimensions (R_x, R_y, R_z) = (0.5,0.5,3) or dot-dashed line, (0.6,0.6,3) or dashed line, and (0.7,0.7,3) or dotted line. B, right plot: Simulations for T^1-universes with dimensions (R_x, R_y, R_z) = (0.5,3,3) or dot-dashed line, (0.6,3,3) or dashed line, and (0.7,3,3) or dotted line. In both pictures the model (R_x, R_y, R_z) = (3,3,3) is represented by a solid line, $S_\circ^{DMR} = 2.59$ (vertical straight line) and the horizontal solid lines indicate the confidence levels of 95%, 90% and 68% (from top to bottom).

In a Euclidean topology the universe is isotropic, and the sum in (3) is normally replaced by an integral. However, in a toroidal universe this is not the case. In this model, only wave numbers that are harmonics of the cell size are allowed. As a result, we have a discrete $\mathbf{k}$ spectrum [6, 15]

$$\mathbf{k} = \frac{2\pi}{R_H}\left(\frac{p_x}{R_x}, \frac{p_y}{R_y}, \frac{p_z}{R_z}\right), \tag{4}$$

where p_x, p_y and p_z are integers.

From equation (3), we can construct simulated skies by calculating [3]

$$\frac{\delta T}{T}(\theta,\phi) \propto \sum_{p_x,p_y,p_z} \left[g_1\cos(2\pi\gamma) + g_2\sin(2\pi\gamma)\right] \alpha^{\frac{n-4}{4}} e^{-(R_H\Theta k_\perp)^2/2} + n(\theta,\phi), \tag{5}$$

where g_1 and g_2 are two independent Gaussian random variables with zero mean and unit variance, $\gamma = \left(\frac{p_x}{R_x}x + \frac{p_y}{R_y}y + \frac{p_z}{R_z}z\right)$, $\alpha \equiv \left(\frac{p_x}{R_x}\right)^2 + \left(\frac{p_y}{R_y}\right)^2 + \left(\frac{p_z}{R_z}\right)^2 \propto k^2$ and n is the spectral index of the scalar perturbations. The term $e^{-(R_H\Theta k_\perp)^2/2}$ represents the experimental beam function, where $k_\perp$ is the length of the $\mathbf{k}$-component perpendicular to the line of sight and Θ is the width of the Gaussian beam given by $\Theta = \mathrm{FWHM}/\sqrt{8\ln 2} \approx 0.43$ FWHM, where FWHM is the full width of the beam at its half maximum. Finally, we model the noise $n(\theta,\phi)$ at each pixel i as independent Gaussian random variables with mean $\langle n_i\rangle = 0$ and variance $\langle n_i n_j\rangle = \sigma_{ij}\delta_{ij}$ [9].

We generate our simulated skies as standard DMR maps with 6144 pixels for n=1, with a Galaxy cut of 20°, FWHM = 7°, the monopole and dipole removed, add noise and normalize to $\sigma_{7^\circ} = 34.98\mu$K (the r.m.s. value at 7° extracted from our DMR map, a 4 year combined 53 plus 90 GHz map with monopole and dipole removed). Fixing a cell size, we construct a simulated sky according to (5), we smooth this once more by 7° and use the statistic defined in (1) to obtain an S-map from which we extract its minimum value $S_\circ$. Repeating this procedure 1000 times, we obtain the probability distribution of $S_\circ$ for that fixed cell size. When we repeat this same procedure for different cell sizes, we are able to construct plots as shown in Figure 2.